

# TISSUE EXPANDER/BREAST RECONSTRUCTION HOME INSTRUCTIONS

## DIET

· Start with a light diet – mainly liquids, then progress to a regular diet.

## ACTIVITY

- · Take it easy and do not schedule any significant activities for the first 4-6 weeks.
- · No lifting more than 5 lbs.
- · Deep breathing exercises should be done every hour while awake for the first 48 hours.

## MEDICATION

- · Do not take aspirin/ibuprofen-type pills unless directed by Dr. Heistein!
- · Take pain medication as prescribed and directed.
- Take Regular or Extra-Strength Tylenol 1-2 tablets every 4-6 hours as needed for lesser pain (but not in combination with the prescribed pain medicine).

## WOUND CARE

- · Do not smoke!
- · Change the dressing daily with sterile gauze pads over the incision and around the drains.
- · Empty the drains every 8-12 hours and record the amount. The drainage bulb must be recompressed after emptying to maintain suction of drainage from the incision.
- · You may have some bloody discharge around the drain tube. You may reinforce the dressing with gauze as needed.
- · Do not shower until 24 hours after the drains are removed.

### PROBLEMS

## Call Dr. Heistein if you have problems or concerns. Observe and call for any of the following:

- · Temperature above 101.
- · Excessive drainage.
- · Increased pain that is not relieved by pain medication.
- · A change in the appearance of your breasts (i.e. if one suddenly appears larger, more painful, or if excessive bruising or hardness is noted).
- · If unable to contact Dr. Heistein in an emergency situation, go to the nearest emergency room.

## ANESTHESIA

· If you had general anesthesia, you may have a sore, dry throat. Throat lozenges and/or gargles may give relief You may have muscle soreness, a jittery feeling, or nausea for 24-48 hours.

## SPECIAL INSTRUCTIONS\_

#### FOLLOW-UP

· Please call (817)820-0000 today to schedule your post-operative appointment for tomorrow (unless previously scheduled)