

BODY LIFT HOME INSTRUCTIONS

DIET

· Start with a light diet – mainly liquids, then progress to a regular diet.

ACTIVITY

- · You must take it easy for 4-6 weeks.
- \cdot Do not do any strenuous activity.
- \cdot Do not try to sit up without using your arms for support.
- · Do not lift more than 5 lbs.
- · Move your legs and feet often while resting.
- · Deep breathe every hour while awake to clear your lungs.

MEDICATION

- · Do not take aspirin/ibuprofen-type pills unless directed by Dr. Heistein!
- · Take pain medication as prescribed and directed.
- · You MUST continue to take the antibiotic until the drains are removed.
- · Take Regular or Extra-Strength Tylenol 1-2 tablets every 4-6 hours as needed for lesser pain (but not in combination with the prescribed pain medicine).

WOUND CARE

- · Do not smoke!
- · Change the dressing daily with dry sterile gauze on the incisions and around the drains also.
- · Wear the abdominal binder at all times except to change the dressing.
- · Empty the drains every 8-12 hours and record the amount. The drainage bulb must be recompressed after emptying to maintain suction of drainage from the incision.
- · You may have some bloody drainage around the drain tube. Reinforce the dressing as needed.
- · Do not shower until 24 hours after the drains are removed.

PROBLEMS

Call Dr. Heistein if you have problems or concerns. Observe and call for any of the following:

- · Temperature above 101.
- · Excessive drainage.
- · Increased pain that is not relieved by pain medication.
- · Heat or redness at or near surgery site.
- · If unable to contact Dr. Heistein in an emergency situation, go to the nearest emergency room.

ANESTHESIA

· If you had general anesthesia, you may have a sore, dry throat. Throat lozenges and/or gargles may give relief. You may have muscle soreness, a jittery feeling, or nausea for 24-48 hours.

SPECIAL INSTRUCTIONS	
FOLLOW-UP	
· Please call tomorrow to schedule your post-operative appointment for	(unless previously scheduled)